

Predicting Multivariate Volatility

Christof Reese and Bernd Rosenow

Institut für Theoretische Physik, Universität zu Köln, D-50923 Germany

(Dated: February 2, 2008)

We suggest two classes of multivariate GARCH-models which are both easy to estimate and perform well in forecasting the covariance matrix of more than one hundred stocks. We apply methods from random matrix theory (RMT) to determine the number of principal components or the number of factors in the multivariate volatility models. In this way only statistically relevant information is used for the estimation of model parameters.

The basic input for the estimation of portfolio risk is a forecast of time varying covariance matrices. Although modern risk management is founded on the analysis of huge data bases and the use of sophisticated theoretical models, there has been only partial progress with respect to covariance forecasts. The reason for this lack of progress is the “curse of dimensionality”: in modern financial engineering, the dimension of the investment universe is comparable to the number of observations, and hence the number of parameters in the covariance or correlation matrix is of the same order as the number of data points.

Traditional multivariate statistics is not able to deal with this kind of problem. Both estimators and tests are consistent only in the limit of L/N going to infinity, where L is the number of observations and N is the dimension of the observed vector [1]. In typical applications, one is dealing with ratios L/N ranging from one to ten, and hence statistical fluctuations due to the lack of a sufficient number of observations dominate an empirically observed covariance matrix. This measurement noise induces randomness in the covariances and hence turns the covariance matrix at least partly in a random matrix.

A covariance matrix can be decomposed into the cross-correlation matrix C and the set of standard deviations $\{\sigma_i^2\}$. To identify the effects of randomness on the eigenvalue spectrum of an empirical cross-correlation matrix, one considers the null hypothesis of completely uncorrelated and identically distributed (i.i.d.) time series. The cross-correlation matrix calculated from these time series is a random matrix R from the so called Wishart ensemble. The eigenvalues of a random Wishart matrix fall in the interval $[\lambda_-, \lambda_+]$ with $\lambda_{\pm} = 1 + N/L \pm 2\sqrt{N/L}$ [2]. This eigenvalue distribution can be compared to that of the empirical cross-correlation matrix C . Agreement between R and C is a sign of randomness, whereas deviations indicate the presence of economically meaningful information. It was found that approximately 94 % of the empirically observed eigenvalues fall in the random matrix interval [3]. The random matrix nature of the main part of the spectrum was proven by comparing the eigenvalue statistics of empirical correlation matrices to the universal predictions of random matrix theory [4]. Only the largest eigenvalues and corresponding eigenvectors carry information about market correlations, busi-

ness sectors and geographical correlations [5, 6]. These economically meaningful correlations are stable in time and hence suitable for forecasting future correlations. Generally, correlations show a higher degree of time stability than covariances which are fluctuating due to time changing volatility.

The time evolution of the variance or volatility is often described with GARCH-models [7], which relate the present variance to past variances and past returns. While GARCH-processes are well suited to describe the dynamics of a univariate volatility process [8], they are less successful in the multivariate setting. In addition to the “curse of dimensionality”, the application of multivariate GARCH-models is hampered by the need to make maximum likelihood estimates of a large number of parameters growing generically like the square of the number of time series. In this note, we suggest two classes of multivariate GARCH-models, which incorporate RMT based progress in the understanding of empirical cross-correlation matrices. These models are both easy to estimate and compare favorably to alternative models as the RiskMetrics covariance estimator [9].

High dimensional correlation matrices

We estimate the sample covariance matrix

$$\Sigma_{ij} = \frac{1}{(L-1)} \sum_{t=1}^L (\epsilon_{i,t} - \langle \epsilon_i \rangle)(\epsilon_{j,t} - \langle \epsilon_j \rangle) \quad (1)$$

and define the sample correlation matrix $C_{ij} = \Sigma_{ij}/(\sigma_i \sigma_j)$ by normalizing with the standard deviations $\sigma_i = \sqrt{\Sigma_{ii}}$. We calculate cross-correlation matrices for the 118 most frequently traded German companies from a) 1711 daily returns $\epsilon_{i,t}$ starting 12/01/93 until 08/31/00, and b) 250 daily returns starting 09/01/99 until 08/31/00, diagonalize them, and rank-order their eigenvalues $\lambda_i < \lambda_k$ for $i < k$. The pdf of the eigenvalues of C_a is displayed in Fig. 1. For both C_a and C_b we find that three eigenvalues lie outside the random matrix interval, but only one of them clearly. The number of eigenvalues clearly outside the RMT-interval is the optimal number of principal components or factors that should be used to forecast the correlation matrix [11].

The largest eigenvalue $\lambda_{\max} = 14.5$ for the seven-year interval is much larger than $\lambda_{\max} = 7.1$ for the one-year interval. This confirms the result of [4, 6, 10] that




FIG. 1: Probability distribution of the eigenvalues of the cross-correlation matrix C^a calculated from daily 118 return time series from 09/01/99 until 08/31/00 (gray line). The bulk of eigenvalues is due to noise and well described by the prediction of random matrix theory (black line). One eigenvalue is clearly separated from the bulk and contains information about market correlations.

the strength of market correlations changes in time. Although the use of time series extending over several years in theory increases the accuracy of correlation estimates, it does not capture the effect of time changing correlations and hence is not advisable.

In order to use only the statistically relevant information in the correlation matrix C , we diagonalize C by an orthogonal transformation X via $X^T C X = \Lambda$. We construct a filtered correlation matrix C'_p by keeping only the p largest eigenvalues in the diagonal matrix [3, 5, 11]

$$\Lambda_{ii}^p = \begin{cases} \Lambda_{ii} & \text{for } i > N - p \\ 0 & \text{for } i \leq N - p \end{cases}, \quad (2)$$

and by transforming back to the original basis $C^p = X \Lambda^p X^T$. For consistency, all diagonal elements of C^p are set to one.

Model description

The basis of the *Sliding Correlation Model* (SCM) is the decomposition of the covariance matrix into the cross-correlation matrix C and standard deviations $\{\sigma_i\}$. Each individual time series is described by a univariate GARCH(1,1) [7] process defined by

$$\sigma_{i,t}^2 = \alpha_{i,0} + \alpha_{i,1} \sigma_{i,t-1}^2 + \beta_{i,1} \epsilon_{i,t-1}^2. \quad (3)$$

Here, $\sigma_{i,t}^2$ and $\epsilon_{i,t}$ are the volatility and the innovation of time series i at time t , respectively. The innovations are modelled as a product $\epsilon_{i,t} = \sigma_{i,t} \cdot x_{i,t}$ with normally distributed $x \in \mathcal{N}(0, 1)$. The parameters $\alpha_{i,0}$, $\alpha_{i,1}$, and $\beta_{i,1}$ are obtained from maximum likelihood estimations. The cross-correlation matrix C is calculated as a moving average to accommodate the change of correlations in time. We use the filtering method described in the last section and keep only the p largest principal components. The covariance matrix in the sliding correlation model SCM(p) of rank p is described by

$$\Sigma_{ij,t} = C_{ij,t}^p \sigma_{i,t} \sigma_{j,t}. \quad (4)$$

This model is related to the constant correlation model [12], where a fixed and unfiltered correlation matrix is used in contrast to our $C_{ij,t}^p$.

Factor Models are motivated by the observation that correlations are large in times of high volatility [6]. In addition, general multivariate GARCH-models can be simplified to factor models, if the parameter matrices do not have full rank. As a special case with parameter matrices of rank one, a one-factor model was discussed in [13]. We suggest a more general model with p factors (abbreviated Factor(p))

$$\Sigma_{ij,t} = \tilde{\sigma}_{i,t}^2 \delta_{ij} + \sum_{k=1}^p \lambda_i^{(k)} \lambda_j^{(k)} \Sigma_t^{(k)}. \quad (5)$$

Here, $\tilde{\sigma}_{i,t}^2$ is the variance of the residual of time series i after linear regression on the p factors $f_t^{(k)}$ with $k = 1 \dots p$, and $\lambda_i^{(k)}$ is the regression coefficient (factor loading) of factor k on time series i . The dynamics of the residuals is described by a GARCH(1,1)-process, and the dynamics of the factors is similarly given by

$$\Sigma_t^{(k)} = \alpha_0^{(k)} + \beta_1^{(k)} \Sigma_{t-1}^{(k)} + \alpha_1^{(k)} (f_{t-1}^{(k)})^2. \quad (6)$$

The factors $f_t^{(k)}$ are defined via an iterative procedure: we use the largest eigenvector $\mathbf{u}^{(N)}$ of C to define $f_t^{(1)} = \sum_{i=1}^N u_i^{(N)} \epsilon_{i,t}$. Next, we perform a linear regression of $f^{(1)}$ on the ϵ_i and recompute the correlation matrix from the residuals. From the new eigenvector corresponding to the largest eigenvalue, we define a factor $f^{(2)}$, perform another linear regression and so on. The number p of factors is chosen according to the number of eigenvalues of the correlation matrix outside the RMT-interval, the time interval for the calculation of factors and factor loadings is not moving but fixed.

The data set for parameter estimation comprises seven years of daily data from 12/01/93 until 08/31/00, which are used for the maximum likelihood estimation of all GARCH parameters, for the calculation of factors and factor loadings of the models Factor(p)-seven, whereas only the last year of daily data is used for models Factor(p)-one. Similarly, the cross-correlation matrix of model SCM(p)-seven is calculated over a sliding time window of length seven year, and a window of length one year is used for model SCM(p)-one.

Our benchmark model is the commonly used Risk Metrics covariance estimator [9] defined by

$$\Sigma_{ij,t} = 0.94 \Sigma_{ij,t-1} + 0.06 \epsilon_{i,t} \epsilon_{j,t}. \quad (7)$$

Test method

We judge the quality of a multivariate volatility model by its ability to forecast the daily variance of both equally distributed and minimum variance portfolios. The predicted portfolio volatility at time t is given by

$$D_{t,\text{predicted}}^2 = \sum_{i,j=1}^N m_{i,t} m_{j,t} \Sigma_{ij,t}. \quad (8)$$




FIG. 2: Prediction of daily volatility of a), b) equally weighted portfolios and c), d) minimum variance portfolios. The black line shows the estimated volatility, the thin grey line the realized volatility calculated from high frequency data while the thick grey line is a seven-day average of the realized volatility. a) Sliding correlation model SCM(1)-seven, which uses one principal component and is estimated over a seven-year period, overestimates the variance by 65 % due to an overestimation of the average correlation strength. b) Model Factor(1)-seven, which uses one factor and is estimated over a seven-year period is able to describe changes in the correlation strength but tends to overestimate correlations in the time after a volatility peak. c) RiskMetrics estimator Eq. 7 predicts a very small variance for a minimum variance portfolio, while the realized variance is a factor 35 larger. d) Sliding correlation model SCM(1)-one uses one principal component and estimates correlations over a sliding time window of one year. The predicted variance is close to the realized one, which is more than 50 % smaller than for the RiskMetrics estimator.

Here, $m_{i,t}$ is the fraction of the capital invested in stock i at time t . For an equally distributed portfolio, we have $m_{i,t} \equiv 1/N$, hence D^2 is just the average element of the covariance matrix. This test probes the ability of a model to correctly predict the average covariance between pairs of stock. On the other hand,

$$m_{i,t} = \sum_{j=1}^N \Sigma_{ij,t}^{-1} / \sum_{k,l=1}^N \Sigma_{kl,t}^{-1} \quad (9)$$

minimize the variance under the constraint that the total invested money is equal to one. The volatility prediction for minimum variance portfolios is a difficult test for volatility models, as the covariance matrix is first used to calculate the portfolio weights and then to estimate the variance of that portfolio. We compare the predicted

volatility to the realized volatility, which is calculated from high frequency (hourly) data [8]. The realized portfolio variance at time t is calculated as

$$D_{t,\text{realized}}^2 = \sum_{i,j=1}^N m_{i,t} m_{j,t} (\langle g_{i,t} g_{j,t} \rangle - \langle g_{i,t} \rangle \langle g_{j,t} \rangle) \quad (10)$$

where the expectation values are taken for 10 hourly returns g_i per trading day. We perform an out of sample test from 09/01/00 until 08/31/01. We calculated the average predicted portfolio variance, the average realized variance, the mean square error (MSE) of the prediction and the mean average percentage error (MAPE).

Test results:

For *equally distributed portfolios*, the performance of the various factor models is very similar to each other. Model Factor(1)-seven, for example, overestimates the overall level of variance but is able to describe the effect of enhanced correlations in high volatility periods (Fig. 2).

The variance estimation of SCM(N)-seven and SCM(1)-seven is very similar to each other, both models severely overestimate the average level of correlations, see Fig. 2 for the results of SCM(1)-seven. The sliding correlation models with a one year time window for correlation estimates provide for the best estimates of realized variances, the RiskMetrics estimator shows comparable results. These results indicate that a good multivariate volatility model must be able to capture not only the change of volatility over time but also the change of the average correlation strength. For details see Table I.

Minimum variance portfolio: Due to the measurement noise contained in its cross-correlation matrix, the model SCM(N)-one clearly underestimates the variance of minimum variance portfolios, the ratio of realized variance to

Model	$D^2_{\text{estimated}}$	D^2_{realized}	MSE	MAPE
SCM(1)-seven	0.654	0.396	0.092	0.886
SCM(2)-seven	0.666	0.396	0.098	0.916
SCM(N)-seven	0.643	0.396	0.086	0.856
SCM(1)-one	0.497	0.396	0.037	0.486
SCM(2)-one	0.500	0.396	0.037	0.493
SCM(N)-one	0.484	0.396	0.034	0.456
Factor(1)-seven	0.568	0.396	0.063	0.615
Factor(2)-seven	0.576	0.396	0.065	0.634
Factor(1)-one	0.571	0.396	0.062	0.618
Factor(2)-one	0.575	0.396	0.064	0.629
RiskMetrics	0.519	0.396	0.057	0.535

TABLE I: Estimation of the variance of equally distributed portfolios

Model	$D^2_{\text{estimated}}$	D^2_{realized}	MSE	MAPE
SCM(1)-seven	0.177	0.243	0.022	0.335
SCM(2)-seven	0.188	0.238	0.025	0.415
SCM(N)-seven	0.190	0.245	0.029	0.409
SCM(1)-one	0.158	0.208	0.016	0.340
SCM(2)-one	0.157	0.208	0.017	0.338
SCM(N)-one	0.128	0.284	0.046	0.479
Factor(1)-seven	0.215	0.304	0.033	0.318
Factor(2)-seven	0.238	0.303	0.035	0.375
Factor(1)-one	0.212	0.300	0.031	0.319
Factor(2)-one	0.220	0.312	0.033	0.305
RiskMetrics	0.013	0.453	0.241	0.967

TABLE II: Estimation of the variance of minimum variance portfolios

predicted variance is 2.22. The filtered model SCM(1)-one performs much better in this respect, the ratio is given by 1.32 (Fig. 2d). In addition, the realized variance of the “filtered” model is 27% lower than that of the original model. Adding one more eigenvalue in the cross-correlation matrix does not change this result much. The performance of the various factor models with one or two factors and estimation periods of seven or one year is quite similar to each other, therefore we discuss the representative model Factor(1)-seven (detailed results can be found in Table II). Although the ratio of realized to predicted variance is with 1.46 better than that of model SCM(N)-one, the realized variance itself is higher than that of the other models. Possibly the performance of the factor models could be increased if the factor loadings were estimated by the maximum likelihood method instead of using the regression coefficients. Details of the results can be found in Table II. All these results compare favorably to the RiskMetrics covariance estimator, which underestimates the portfolio variance by a factor of 35 (Fig. 2c). Also, the results reported in [14] for DCC-GARCH (overestimation of risk by several hundred percent) are much less precise than the estimates of SCM(1)-one, for example.

In summary, we have shown that the “curse of dimensionality” can be overcome by using appropriate statistical tools relying on RMT. Building on the distinction between statistically relevant information and noise in the correlation matrix, we have suggested two types of multivariate GARCH-models that are easy to estimate even for more than 100 time series. Both the sliding correlation model and the factor model studied have a much better prediction quality than the industry standard RiskMetrics covariance estimator.

-
- [1] R. B. Muirhead, *Aspects of Multivariate Statistical Theory*, Wiley New York (1982).
 - [2] V.A. Marčenko and L.A. Pastur, *Math. U.S.S.R. Sbornik* **1**, 457 (1967).
 - [3] L. Laloux, P. Cizeau, J.-P. Bouchaud, and M. Potters, *Phys. Rev. Lett.* **83**, 1467; see also, L. Laloux et al., *Int. J. Theor. Appl. Finance* **3**, 391 (2000). (1999).
 - [4] V. Plerou, P. Gopikrishnan, B. Rosenow, L.A.N. Amaral, and H.E. Stanley, *Phys. Rev. Lett.* **83**, 1471 (1999).
 - [5] P. Gopikrishnan, B. Rosenow, V. Plerou, and H.E. Stanley, *Phys. Rev. B* **64**, 035106R (2001); see also, P. Gopikrishnan et al., preprint cond-mat/0011145.
 - [6] V. Plerou, P. Gopikrishnan, B. Rosenow, L.A.N. Amaral, T. Guhr, and H.E. Stanley, *Phys. Rev. E* **65**, 066136 (2002).
 - [7] R.F. Engle (Ed.), *ARCH Selected Readings*, Oxford University Press (1995).
 - [8] T. Anderson and T. Bollerslev, Technical Report 6023, National Bureau of Economic Research (1997).
 - [9] J. Longestaey, P. Zangari, *RiskMetrics-Technical Document*, 4th ed., New York, Morgan Guaranty Trust Co.
 - [10] S. Drozd, F. Gruemmer, F. Ruf, and J. Speth, *Physica A* **287**, 440 (2000).
 - [11] B. Rosenow, P. Gopikrishnan, V. Plerou, and H. E. Stanley, *Portfolio optimization and the random magnet problem*, *Europhys. Lett.* **59**, 500 (2002).
 - [12] T. Bollerslev, *Review of Economics and Statistics* **72**, 498 (1990).
 - [13] R. Engle, V. Ng, M. Rothschild, *J. of Econometrics* **45**, 213 (1990).
 - [14] R. Engle, K. Sheppard, *Theoretical and Empirical properties of Dynamic Conditional Correlation Multivariate GARCH*, Research Paper (2001).